

Cancer and blood clots

What you should know when
you are in hospital

Information for patients



Leeds Cancer
Centre

Please read this leaflet carefully. It will give you information about decreasing the chance of developing a blood clot during and after your hospital stay.

If you are living with cancer you will be aware that both the cancer and its treatment are associated with potential complications. One such complication is an increased risk of developing blood clots (also known as venous thromboembolism).


What is Venous Thromboembolism (VTE)?

There are two types of VTE:

- **Deep Vein Thrombosis (DVT):** a DVT is a blood clot (also known as a thrombosis) that forms in a deep vein, most commonly in the legs but they can also occur in the arms.
- **Pulmonary Embolism (PE):** if all or part of the DVT breaks free and passes through your blood vessels, it can reach your lungs. This is called a PE.




The vein in the leg



Normal blood flow



A clot has formed




Part of the clot has broken off

Signs and symptoms of VTE

DVT - Symptoms can include swelling, redness/ discolouration, warmth and tenderness/pain that may be worse when standing or walking. Occasionally there are no symptoms except pain.

PE - Symptoms can include coughing, chest pain, breathlessness, blood stained phlegm and collapse.



If you develop any of these symptoms please get medical advice immediately.

Cancer Associated Thrombosis (CAT) is the name for a DVT or PE that develops when you also have cancer. Unfortunately, VTE is common in people with cancer.

There are several reasons why you might develop a VTE when you have cancer, they are detailed below:

Type of Cancer

Having active cancer can increase your risk of developing a VTE.

The types of cancer that are most likely to increase your risk of VTE are detailed below:

- Stomach
- Pancreas
- Lung
- Lymphoma
- Gynaecological
- Brain
- Kidney

Cancer treatments

Some cancer treatments can increase your risk of developing a VTE. The types of treatment that are most likely to increase your risk of VTE are detailed below:

- Chemotherapy - particularly cisplatin
- Hormone therapy such as: tamoxifen and anastrozole
- Immunomodulatory drugs such as: thalidomide and lenalidomide
- Antiangiogenic therapies such as bevacizumab and sunitinib
- Supportive agents such as erythropoietin and darbopoetin.

Other effects

Reduced mobility due to cancer treatments, blood vessel damage from surgery, or having a central venous catheter are other possible risk factors for getting a VTE.

Other risk factors for VTE

- Past history of blood VTE's
- Being overweight
- Reduced mobility (not walking or moving around)
- Combined contraceptives (i.e. the Pill, the contraceptive patch (Evra®) or the vaginal ring (Nuvaring®))
- Hormone replacement therapy (HRT)
- Recent surgery especially on hips or knees
- Age over 60 years
- Pregnancy or given birth in the past six weeks

- Dehydration
- Smoking
- Varicose veins with phlebitis
- A close relative has had a blood VTE e.g. mother, father, brother, sister, child
- Known thrombophilia (inherited blood clotting disorders)
- Long distance travel of more than four hours within the last eight weeks.

Fortunately, by becoming aware of the signs and symptoms of VTE and always keeping your healthcare providers informed, you can reduce the risks associated with this condition.


How can I help to reduce my risk of developing a VTE?

You can help reduce the risk of developing a VTE by:

- keep drinking plenty of fluids (unless advised otherwise);
- moving around as much as you can;
- wearing the anti-embolism stockings or other devices you are given;
- taking any medication you have been prescribed.

Carry out the following leg exercises (even if you are in bed):

- Lying on your back or sitting, bend and straighten your ankles quickly 10 times.
- Keep your knees straight during the exercise to stretch your calf muscles.
- Try to do this exercise two or three times an hour.



How will hospital staff help me reduce my risk while I am in hospital?

There are a number of methods that can be used to reduce your risk of developing a VTE whilst you are in hospital. The methods that you use during your hospital stay will depend on your level of risk.

When you come into hospital your doctors will assess your risk of developing a VTE. During your hospital stay your risk may change so you will be re-assessed throughout your stay.

Some of the ways that we reduce the risk are:

- Encouraging you to remain as mobile as possible. This may include asking you to regularly perform the leg exercises discussed on the previous page.
- Encouraging you to drink plenty of fluids. If you cannot drink, this fluid may be provided by a drip into your vein.
- Offering you elastic stockings to be worn on both legs (these are usually worn after surgery). These encourage blood flow through the legs and reduce the risk of VTE's. It is important that stockings are the correct size so your legs will be measured to ensure correct fit. You may find them a little uncomfortable and tight at first, but most people get used to them.



- If you are in bed for a long time then you may be fitted with pumping devices that are wrapped around each of your legs. They are very simple to use and feel similar to a blood pressure cuff.



The device is inflated on your legs regularly for short periods to encourage blood flow.

- Giving you a daily injection of a low dose heparin. This medicine thins the blood and makes VTE's less likely to form. These injections go just under the skin into the fatty layer and are usually given in the stomach, thigh or upper arm.



The injection is usually given at approximately 6pm in the evening.

You may need to continue these injections for a little while after you have been discharged home so the nursing staff will teach you how to administer the injection yourself or a relative/friend can be taught how to administer them for you. If you do not think you will be able to administer the



injections yourself please think about who you would like to do it for you. They will need to come into hospital when you are due to have your injection so the nursing staff can teach them how to administer it for you.

Low dose heparin is made from pork derived heparin sodium. If you have any questions or concerns about this please speak to your nurse, pharmacist or doctor.

There is a risk of bleeding following use of heparin. You will be assessed for your risk of bleeding. Let the doctors know if you develop unexplained bruises or any bleeding.

If you have any questions about the risk of developing a VTE whilst in hospital or about any of the things described above to reduce the risk, speak to your doctor or nurse.

How do I use the elastic stockings?

To ensure the stockings are effective:

- Ensure the stockings are not wrinkled. Otherwise they will dig into the skin underneath and can cause tissue damage.
- Do not turn/roll down the top of the stockings. This may interfere with the blood flow in your legs.
- If you experience tingling or numbness in your legs tell the nurses. This means that the stockings are too tight. The nurse will need to reassess your legs.
- The stockings should be worn day and night while you are immobile.
- You or your nurse should remove the stockings at least once a day to wash and dry your legs. Apply a non-perfumed moisturiser after you dry your legs. Whilst washing your legs look at the skin. If there are any signs of blisters or red marks that do not disappear, particularly on the heels, shin and toes, you should stop wearing the stockings. Tell a nurse if you see these marks.

How will I be monitored during my admission?

Members of the nursing and medical team will review your progress throughout your in-patient stay but it is important to report to the nurse or doctor any of the following symptoms:

- Breathlessness
- Chest pain
- Tenderness, swelling or pain in your calf area.
- Bleeding or unexplained bruising (if you are being given heparin injections).

How can I reduce my risk after going home from hospital?

You should continue to drink plenty of water or other non-alcoholic drinks to keep hydrated (unless advised otherwise) and move around as much as you can.

If you are at high risk of developing a VTE you will need to continue to reduce your risks when you go home.

You may need to wear stockings for a period of time at home or be given some exercises to do at home. You may be asked to continue low-dose heparin injections to thin your blood for some time after you go home. If this is the case your doctor or nurse will talk to you about this before you go home.

What do I do if I develop symptoms when I go home?

If you develop sudden shortness of breath and or chest pain:


Ring 999

If you develop new swelling, tenderness and/or pain of your leg or arm you should ring:

0113 243 3144 and ask for the Oncology Patient Enquiries Bleepholder

Useful sources of information

- Please ask your doctor, nurse or pharmacist for more information.
- NHS choices website for patient information on blood clots. Visit www.nhs.uk/Conditions/Thrombosis
- Lifeblood: The thrombosis charity also has more information. Visit www.thrombosis-charity.org.uk
- Anticoagulation Europe for information on anticoagulation. Visit www.anticoagulationeurope.org



What did you think of your care?

Scan the QR code or visit bit.ly/nhsleedsfft

Your views matter



The printing of this leaflet has been
funded by Leeds Hospitals Charity
leedshospitalscharity.org.uk

leeds hospitals charity

charity number: 1170369

© The Leeds Teaching Hospitals NHS Trust • 2nd edition Ver 1
Developed by: Louise O'Prey, VTE Prevention Nurse
Produced by: Medical Illustration Services • MID code: 20220210_008/JG

LN003916
Publication date
04/2022
Review date
04/2024