



**The Leeds
Teaching Hospitals**
NHS Trust

Endoscopy Unit

Propofol sedation for Endoscopic procedures

Information for
patients

This leaflet has been designed to give you information about receiving Propofol sedation, while having your Endoscopic procedure. You will be given separate information about the procedure you are having and how to prepare for your procedure.

What is propofol sedation?

Propofol sedation is a form of deep sedation administered by an anaesthetist, a qualified doctor who has had specialist training to care for you before, during and after any procedures. The level of sedation can be increased or decreased, as required, by the anaesthetist to enable adequate sedation to be maintained throughout the procedure.

Propofol sedation is effective and safe, it is the modern alternative to general anaesthesia for endoscopy procedures, which wears off very quickly without a 'hangover effect'.

Although propofol sedation is different to general anaesthesia, often, patients cannot tell the difference between the two as they usually remember nothing of the procedure.

There are a small number of complex procedures or pre-existing health conditions where your anaesthetist may feel a general anaesthetic is more suitable, you will be advised of this before the procedure.

If anything in this leaflet worries you or you have questions that we have not answered, please discuss with your anaesthetist or the specialist nurse during your pre-operative assessment telephone call.

Propofol sedation advantage?

- Compared to other sedation techniques: adjusted to individual needs and infusion continues until procedure complete.
- Patient feedback: "it was 1,000 times better than last time".
- Compared to General anaesthesia: quicker recovery, less drops in blood pressure and you will position yourself comfortably before being sedated.
- Patient feedback: "I don't remember anything", "No pain, not feeling ill, no sickness, feeling good overall".

Possible risks of having Propofol sedation

- Your breathing rate may become slow. This is common with deep sedation but is a risk whenever sedation is used. The anaesthetist will be monitoring you at all times and will take the appropriate action, if needed.
- Occasionally, your blood pressure may drop but again, the anaesthetist will be monitoring this.
- An allergic reaction to the sedation is very rare.
- You may have a small bruise where your cannula was placed or attempted.
- After the procedure, you may feel less steady on your feet and be at higher risk of falling, especially if you are elderly or have existing mobility problems.

Pre-operative assessment

A member of the specialist nursing team will contact you by telephone before the day of the procedure.

They will ask you a lot of questions about your general health and the medications you are currently taking.

The team will also need to know how you will get home and who is going to look after you after the procedure. You must have a responsible adult with you for 12 hours after the procedure.

The specialist nurse will also ask for your height and weight so please can you make sure you have this information readily available.

On the day of the procedure

Do not eat for at least six hours before your appointment time. Sometimes, this may be longer (you will be advised of this in your appointment letter) if you need to prepare your bowel for the procedure. You may drink water up to two hours before your appointment time. This is to make sure that you keep hydrated before your procedure.

Please take all of your usual medications on the morning of your procedure unless the specialist nurse has advised you otherwise. Bring all your medication, including inhalers into hospital with you.

You should also bring warm clothing such as a cardigan or jumper. Our hospital is air-conditioned and it is important that you do not get cold while waiting for your procedure.

What will happen before my procedure?

You will come into the Endoscopy Department where you will be seen by the anaesthetist before the procedure to discuss any questions you may have.

You will have a cannula (drip line) inserted in the back of your hand, either in the Endoscopy Department, or in the procedure room. You may need to change into a hospital gown.

When you go for your procedure, we will attach our routine monitoring equipment and give you oxygen via small tubes that fit comfortably into your nostrils.

Depending on your procedure, the anaesthetist may spray some local anaesthetic into the back of your mouth and ask you to swallow it. This will numb your throat but you will be able to breathe normally.

How is the sedation given?

We will attach your drip through the cannula in the back of your hand, which the sedation is administered through and then ask you to lie down, usually on your side or front.

We will then start the sedation medication. The medication may feel cold or slightly stinging at first but this will quickly go away.

The anaesthetist will be with you at all times watching you closely to make sure you are at the right level of sedation.

The procedure will not start until you are deeply sedated. At the end of the procedure, the sedation is stopped and you will wake up quickly.

Recovery

You will return to the Endoscopy department where you will be observed. If you have had the throat spray, you will not be able to eat or drink anything for one hour after the time the throat spray was given. This is to allow the numbness to wear off. Once you have had something to drink and you are stable, you will be able to go home.

Going home

Someone will need to collect you and take you home. You must also have someone stay with you at home for 12 hours. **You should not drive, drink alcohol or use heavy machinery for 24 hours after your sedation.**

Further resources

Sedation explained by the Royal College of Anaesthetists

https://rcoa.ac.uk/sites/default/files/documents/2021-07/12-SedationExp2021_0.pdf



Contact information

In the unlikely event that you have any problems after discharge, you can call the Endoscopy department for help and advice or contact your GP.

Endoscopy department

Telephone: **0113 206 7833**

Monday - Friday, 7.30 am - 4.30 pm

(outside of these hours, please contact NHS 111 for advice)



What did you think of your care?

Scan the QR code or visit bit.ly/nhsleedsfft

Your views matter



© The Leeds Teaching Hospitals NHS Trust • 1st edition (Ver 1)
Developed by: Dr Alison Colhoun, Consultant (Anaesthetics) and
Dr Nilmini Manawaduge, Locum Consultant (Anaesthetics)
Produced by: Medical Illustration Services • MID code: 20230717_004/IH

LN005662
Publication date
03/2024
Review date
03/2027